

Abstract Submitted  
for the GEC19 Meeting of  
The American Physical Society

Sorting Category: 02.01 (C)

**Electron kinetics in fast-pulsed discharges**<sup>1</sup> LL ALVES, A TEJERO-DEL-CAZ, V GUERRA, N PINHÃO, CD PINTASSILGO, T SILVA, IPFN/IST-UL — This work presents a comparative study of the electron kinetics in fast-pulsed discharges produced in dry air (80% N<sub>2</sub> - 20% O<sub>2</sub>), considering a quasi-stationary and a time-dependent solution of the electron Boltzmann equation (EBE), written under the classical two-term approximation. The simulations were performed using the open-source LisbOn KInetics Boltzmann solver (LoKI-B) [1,2], which handles simulations in any atomic / molecular gas mixture, considering first and second kind collisions with any target state (electronic, vibrational and rotational), characterized by any user-prescribed population. The original capabilities of LoKI-B have been extended in order to obtain the time-dependent solution of the EBE for a pulsed electric field. Noticeable deviations, between the quasi-stationary and the time-dependent calculations of the electron energy distribution function and related macroscopic coefficients, are observed below the microsecond scale, for a pulse peaking at 45 Td with rise and fall times of  $\sim 1\mu\text{s}$  and  $\sim 10\mu\text{s}$ , respectively.

[1] A. Tejero-del-Caz *et al*, Plasmas Sources Sci. Technol. **28** (2019) 043001

[2] <https://github.com/IST-Lisbon/LoKI>

<sup>1</sup>This work was funded by Portuguese FCT Fundação para a Ciência e a Tecnologia, under project UID/FIS/50010/2019.

Prefer Oral Session  
 Prefer Poster Session

Luis Alves  
llalves@tecnico.ulisboa.pt  
IPFN/IST-UL

Date submitted: 30 May 2019

Electronic form version 1.4